arXiv:physics/0302102v1 [physics.ed-ph] 28 Feb 2003

A NOTE ON DERIVATION OF RUTHERFORD FORMULA WITHIN BORN APPROXIMATION

Michal Demetrian*
Department of Theoretical Physics
Faculty of Mathematics, Physics and Informatics
Mlynska Dolina F2, 842 48 Bratislava IV
Slovak Republic

February 2, 2008

Abstract

It is shown in this paper that one does not need to use just exponential dumping factor when computing the Rutherford formula within Born approximation. Text, which is very simple, might be of interest for physics students as well as for physics teachers.

Example 1

The scattering amplitude for a particle of mass m in the spherically symmetric potential energy $V(r)$ within the Born approximation is given by

$$f(\theta) = -\frac{2m}{\hbar^2\kappa}\int_0^\infty \mathrm{d}r r V(r)\sin(\kappa r) \quad , \tag{1}$$

where $\vec{\kappa} = \vec{q} - \vec{q}'$, $\vec{q}$ is the wave vector of incident particle and $\vec{q}'$ is the wave vector of scattered particle. The vector κ is related to the scattering axial angle θ by the equation $\kappa = 2q\sin(\theta/2)$. If we insert the Coulomb law

$$V(r) = \frac{e^2}{4\pi\epsilon_0 r}$$

into (1) instead of V we get

$$f(\theta) = -\frac{2m}{\hbar^2\kappa}\frac{e^2}{4\pi\epsilon_0}\int_0^\infty \mathrm{d}r\sin(\kappa r) \quad .$$

The above written integral does not converge, of course. The standard procedure to make the integral to have a sense is to regularize the Coulomb law by multiplying it by the function $\exp(-\mu r)$, where $\mu > 0$ and after having performed

---

* *demetrian@fmph.uniba.sk*

integration to make the limit $\mu \to 0^+$. We will show that the regularizator has not to be of the form $\exp(-\mu r)$.

Let $\{g_n(r)\}_{n=1}^{\infty}$ be the sequence of monotonically decreasing functions defined on the half-line $r \geq 0$ such that $\{g_n\}$ converges point-wise to 1 and we request that $g_n(0) = 1$ and $\lim_{r\to\infty} g_n(r) = 0$. Moreover we request that the derivatives $g_n'(r)$ approach zero at $r \to \infty$ uniformly with respect to n.
The scattering amplitude (1) for the potential energy

$$V_n(r) = \frac{e^2}{4\pi\epsilon_0}\frac{1}{r}g_n(r)$$

is given by

$$f_n(\theta) = -\frac{2m}{\hbar^2\kappa}\frac{e^2}{4\pi\epsilon_0}\int_0^\infty \mathrm{d}r\sin(\kappa r)g_n(r) \equiv -\frac{2m}{\hbar^2\kappa}\frac{e^2}{4\pi\epsilon_0}I_n(\kappa) \quad .$$

The integral $I_n(\kappa)$ exists (as Riemann integral) due to our assumptions on the functions g_n and the Dirichlet's criterion. Let us compute the integral $I_n(\kappa)$ *per partes.* We get

$$I_n(\kappa) = \frac{1}{\kappa} + \frac{1}{\kappa}\int_0^\infty \mathrm{d}r\cos(\kappa r)g_n'(r) \quad .$$

Our task is to show that the second term goes to zero as n goes to infinity. But this is true because of the properties of the sequence $\{g_n\}$ and because of the fact that there exists such positive constant $K > 0$ that for all $A > 0$ we have

$$\left|\int_0^A \mathrm{d}x\cos(\kappa x)\right| \leq K \quad .$$

Therefore we have

$$\lim_{n\to\infty} f_n(\theta) = -\frac{2m}{\hbar^2}\frac{e^2}{4\pi\epsilon_0}\frac{1}{\kappa^2} \quad ,$$

from which we get the differential cross-section

$$\frac{d\sigma}{\mathrm{d}\Omega} = |f_n(\theta)|^2 = \frac{4m^2e^4}{\hbar^4(4\pi\epsilon_0)^2\kappa^4}$$

which is nothing else but the Rutherford formula.
As an example we can take the sequence of functions

$$g_n(r) = 1 - \frac{2}{\pi}\arctan\left(\frac{r}{n}\right) \quad .$$

In this case we are able to compute $f_n(\theta)$ in terms of elementary functions. In fact:

$$\begin{aligned} I_n(\kappa) &= \frac{1}{\kappa} - \frac{1}{\kappa}\frac{2}{n\pi}\int_0^\infty \mathrm{d}r\frac{\cos(\kappa r)}{1+\left(\frac{r}{n}\right)^2} = \frac{1}{\kappa} - \frac{1}{\kappa}\frac{1}{\pi}\int_{-\infty}^\infty \mathrm{d}x\frac{\exp(i\kappa n x)}{1+x^2} \\ &= \frac{1}{\kappa}\left[1 - 2\pi i\lim_{z\to i}(z-i)\frac{\exp(i\kappa n z)}{1+z^2}\right] = \frac{1}{\kappa}\left[1-\exp(-\kappa n)\right] \to^{n\to\infty} \frac{1}{\kappa} \quad . \end{aligned}$$

We mention, that it is possible to take less limiting assumption on g_n to get the same result.

Example 2

In this section we will show another possibility how to regularize the Coulomb potential. Now, let us take the potential

$$V(r,a) = \frac{e^2}{4\pi\epsilon_0}\frac{1}{r^{1+a}}, \quad 0 < a < 2 \quad . \tag{2}$$

Then the scattering amplitude has the form

$$f(\theta,a) = -\frac{2m}{\hbar^2\kappa}\frac{e^2}{4\pi\epsilon_0}\int_0^\infty \mathrm{d}r\frac{\sin(\kappa r)}{r^a} \quad . \tag{3}$$

So, we have to compute the following integral

$$I(a,\kappa) = \int_0^\infty \mathrm{d}r\frac{\sin(\kappa r)}{r^a} \quad . \tag{4}$$

To do this it would be useful to realise that the following identity holds

$$\frac{1}{x^a} = \frac{1}{\Gamma(a)}\int_0^\infty \mathrm{d}t t^{a-1}e^{-tx} \quad . \tag{5}$$

Inserting (5) into (4) we have

$$\begin{aligned} I(a,\kappa) &= \frac{1}{\Gamma(a)}\int_0^\infty \mathrm{d}t t^{a-1}\int_0^\infty \mathrm{d}r\sin(\kappa r)e^{-tr} = \frac{\kappa}{\Gamma(a)}\int_0^\infty \mathrm{d}t\frac{t^{a-1}}{\kappa^2+t^2} \\ &= |t=\kappa u| = \frac{\kappa^{a-1}}{\Gamma(a)}\int_0^\infty \mathrm{d}u\frac{u^{a-1}}{1+u^2} = \left|\frac{1}{1+u^2} = w\right| \\ &= \frac{\kappa^{a-1}}{2\Gamma(a)}\int_0^1 \mathrm{d}w(1-w)^{a/2-1}w^{a/2} = \frac{\kappa^{a-1}}{2\Gamma(a)}\frac{\pi}{\sin\left(\frac{\pi a}{2}\right)} \quad . \end{aligned}$$

So, the scattering amplitude for the potential (2) is given by the following formula

$$f(\theta,a) = -\frac{2me^2}{4\pi\epsilon_0\hbar^2\kappa^{2-a}}\frac{\pi}{2\Gamma(a)\sin\left(\frac{\pi a}{2}\right)} \quad , \tag{6}$$

from which we get in the limit $a \to 0^+$ the amplitude

$$f(\theta) = -\frac{2me^2}{4\pi\epsilon_0\hbar^2\kappa^2}$$

which leads just to the Rutherford formula.